

PERMANENT GENETIC RESOURCES NOTE

Novel primers for the amplification of nuclear DNA introns in the entomopathogenic nematode *Heterorhabditis bacteriophora* and their cross-amplification in seven other *Heterorhabditis* species

S. O. REGEAI,*† K. M. DOLAN,*‡ D. A. FITZPATRICK,§ J. A. BROWNE,*¶ J. T. JONES** and A. M. BURNELL*

*Biology Department, National University of Ireland Maynooth Co. Kildare, Ireland, †Faculty of Science, Al-Fatah University, PO Box 13618, Tripoli, Libya, ‡Applied Biosystems, Lingley House, 120 Birchwood Boulevard, Warrington, Cheshire WA3 7QH, UK, §School of Biomolecular and Biomedical Science, Conway Institute, ¶Veterinary Sciences Centre, University College Dublin, Belfield, Dublin 4, Ireland, **PPP Programme, SCRI, Invergowrie, Dundee DD2 5DA, UK

Abstract

We describe 24 novel primers that amplify intron regions in housekeeping and structural genes of *Heterorhabditis bacteriophora*. The cross-amplification potential of these primers in seven other *Heterorhabditis* species was determined. The results obtained showed inter-specific nucleotide, length and splice site variability in the sequenced introns and for one gene, an intron gain was observed. These primers will be useful tools for studying population genetics, genetic diversity and intron DNA evolution within the genus *Heterorhabditis* and other genera of rhabditid nematodes.

Keywords: entomopathogenic, EPIC-PCR, *Heterorhabditis*, intron, molecular markers, nematode

Received 9 May 2008; revision 12 May 2008

Heterorhabditid nematodes are obligate and lethal parasites of insects. The infective juvenile (IJ) stages of *Heterorhabditis* are modified third-stage juveniles which occur in soil and are responsible for locating and infecting an insect host. On gaining entry to the insect haemocoel, IJs release cells of a bacterial symbiont of the genus *Photorhabdus* which they harbour in their intestine (Boemare *et al.* 1993). The growth of *Photorhabdus* cells in the infected insect provides a suitable medium for nematode growth and reproduction. Each *Heterorhabditis* IJ matures to become a self-fertile hermaphrodite female, but the progeny of this female contains amphimictic males and females (Dix *et al.* 1992). Approximately 2 weeks post-infection, up to half a million IJs per gram of insect are produced (Akhurst & Bedding 1986). These IJs emerge into the soil where they may survive for several months in the absence of a suitable host. Since a single hermaphrodite IJ can establish a population that undergoes explosive expansion in an insect cadaver, populations of *H. bacteriophora* may be subjected to severe

genetic bottleneck effects, but whether these founder effects lead to spatially distinct populations or whether migration between local populations maintains population cohesiveness has not been determined. Entomopathogenic nematodes (EPN) have been studied intensively because of their applications in biological control. Several studies have been published on the molecular phylogeny and molecular diagnostics of these nematodes (reviewed by Adams *et al.* 2006), but studies on the population genetics of EPN are lacking.

We developed novel primer sets for the amplification of introns from 24 structural and housekeeping genes from *H. bacteriophora*, and we tested the ability of these primers to amplify homologous introns in seven additional species of *Heterorhabditis*. Introns harbour high levels of sequence and length polymorphisms (Palumbi & Baker 1994; Graur & Li 2000). Thus, introns are suitable molecular markers for studies of population structure within and among species, and also for reconstructing relationships among closely related species. We constructed a cDNA library from IJs of *H. bacteriophora* (strain HP88) in the vector pBluescript (SK+) (Stratagene) in *Escherichia coli* DH10B cells. Clones

Correspondence: Ann M. Burnell, Fax: +353 17083840; E-mail: ann.burnell@nuim.ie.




Fig. 1 Agarose gel showing the sizes of gDNA and cDNA amplicons from *Heterorhabditis bacteriophora* (HP88 strain). A larger gDNA amplification product indicates the presence of introns in the gDNA sequence. M (100 bp DNA ladder). Each pair of numbered lanes shows, respectively, gDNA and cDNA amplification products for a single pair of PCR primers. The PCR primers were as follows: 1–2, F16H9.1a (*rgs-2*); 3–4, R11A5.4 (*pepck*); 5–6, M106.5 (*cap-2*); 7–8, T20G5.1 (*chc-1*) (F1/R1); 9–10, C09B8.1 (*ipp-5*); 11–12, F08B6.4b (*unc-87*); 13–14, T20G5.1 (*chc-1*) (F2/R2), bands of equal size indicating absence of an intron; 15–16, F11C3.3 (*unc-54*); 17–18, Y54G11 A5a (*cat*). The primer details are presented in Table 1.

were randomly sequenced from this library, and from these expressed sequence tag (EST) sequences, data for 24 conserved housekeeping and structural genes were used to design exon-primed intron crossing (EPIC) polymerase chain reaction (PCR) primers (Palumbi & Baker 1994).

Putative intron positions in *H. bacteriophora* genes were identified by aligning homologous cDNA and gDNA sequences from *Caenorhabditis elegans* and *C. briggsae* with EST sequences from *H. bacteriophora* and by locating characteristic splice site motifs at the 5' (e.g. GT) and the 3' (e.g. AG) ends of these putative intron sites. PCR primers were designed manually to anneal to *H. bacteriophora* sequences in the flanking exons 30–80 base pairs from each putative intron. For each primer pair, the target sequence was amplified in both gDNA and cDNA from *H. bacteriophora* HP88 strain to detect size differences in the PCR product between the two templates (Fig. 1). PCR amplifications were performed with 0.25 μ M of each primer, 200 μ M of each dNTP, 1 mM MgCl₂, 1 \times reaction buffer, and 1 U of *Taq* DNA polymerase (Promega), in a final volume of 25 μ L. Cycling conditions were 94 $^{\circ}$ C for 1 min, 55–62 $^{\circ}$ C for 1 min (the annealing temperature ranged from 55 to 62 $^{\circ}$ C depending on primers used), 72 $^{\circ}$ C for 1 min, for 30 cycles, and a final extension at 72 $^{\circ}$ C for 10 min. PCR products were visualized in ethidium bromide-stained 1% agarose gels. When putative introns were detected in the gDNA template, the PCR was repeated using High Fidelity Platinum *Taq* DNA polymerase (Invitrogen). PCR products were purified using a QIAGEN PCR Purification Kit. The purified products were cloned into TOPO TA cloning kit (Invitrogen) and the inserts were sequenced (MWG-BIOTECH).

Twenty-four primer pairs were designed (Table 1) and their cross-amplification in seven other *Heterorhabditis* species was also determined (Table S1, Supporting Information). Two primers (C24F3.2 and C04F6.3) did not amplify in species other than *H. bacteriophora* and another four primers (Y54G11 A5a, M106.5, T20G5.1 and *odr-3*) were tested in *H. bacteriophora* only. A total of 169 introns

from eight *Heterorhabditis* species were identified and analysed (Table S2). The *H. bacteriophora* intron borders were determined by aligning the sequences of the *H. bacteriophora* cDNA (EST) and gDNA EPIC-PCR products. The intron borders in the genomic sequences of the seven additional *Heterorhabditis* species were inferred by aligning the gDNA EPIC-PCR products from these species with the homologous *H. bacteriophora* EPIC-PCR sequence in which the intron border had already been determined. As additional checks, the flanking exon sequences were concatenated and translated (using Translate Tool at <http://expasy.org/>). The resulting amino acid sequences were subjected to BLAST analysis against the GenBank (<http://www.ncbi.nlm.nih.gov>) and Wormbase (<http://www.wormbase.org>) databases to confirm gene identity and the correct reading frame. The amino acid translations of the flanking exon sequences of all eight species of *Heterorhabditis* were also aligned with each other and with the homologous sequence from *C. elegans* using Clustal_X (Thompson *et al.* 1997).

The results of EPIC-PCR among *Heterorhabditis* species showed interspecific nucleotide, length and splice site variability in the sequenced introns, and for one gene, an intron gain was observed. These novel intron primers provide molecular markers which will serve as useful tools for studying the population genetics and genetic diversity and intron DNA evolution within the genus *Heterorhabditis*. Because these primers have been identified and designed utilizing their homology to conserved housekeeping and structural gene sequences in *C. elegans*, it is likely that many of these could also be used in studies of other nematode species for which DNA sequence data are currently not available.

Acknowledgements

This research was funded by Enterprise Ireland and the Libyan Ministry of Higher Education. SCRI receives funding from the Rural and Environment Research and Analysis Directorate of the Scottish Government.

Table 1 EPIC-PCR primers used in this study, PCR product sizes and intron lengths in *Heterorhabditis bacteriophora*

Gene name/intron no.	GenBank Accession no.	Primer name	Primer sequence 5'-3'	PCR product size (bp)	Intron size (bp)
ADP-ribosylation factor 1 (<i>arf-1</i>) B0336.2, intron 1	EE724169	arf1F arf1R	CTGATGGTAGGACTAGATGC CCCTAAGTTCATCTTCAGCC	483†	76 115
Dual specificity phosphatase C24F3.2, intron 1	EE724178	dspF dspR	GGTGCAAGTTCAGTTATCAGC GCCTGATAGTTGAGACTTTTCG	798‡	205 118 63
Catalase (Y54G11A.5a)	EE724180	catF catR	CATCGTGATGGACCGGCTTGC CACACCACGGTTCTCCTGGG	220§	100
Myosin heavy chain class II (<i>unc-54</i>) F11C3.3, intron 6	EE724181	unc54F unc54R	GCTCAAGCTGGTTCAGTACGAGC ACTGCGCTCTTCGGCAGCC	190	76
Zinc finger protein (K02D7.2), Intron 2	EE724182	zinc FPF zinc FPR	CCAAACGTGTGCGCACTGC CACACTGCTCACAGATACAAGG	723	625
Inositol polyphosphate 5-phosphatase (<i>ipp-5</i>) C09B8.1, intron 5	EE724183	ipp5F ipp5R	GAGCAGCTATCCGTACAGAG CGAGAGGATGGGGTCCGG	367	240
Clathrin heavy chain T20G5.1 (<i>chc-1</i>)	EE724184	chc1F1 chc1R1 chc1F2 chc1R2	CGATTGGAACGAGCAGAAGCAG CCATACCTGGACCAGGAGC GGACACCGCAGTATAGTGCCC CCATGCAAGATCGAACCACAAAC	200§	150 180 0
F-actin capping protein (<i>cap-2</i>) M106.5	EE724185	cap2F cap2R	CTCCGACGTCTAGAAATCGAAGC GGACGATACTCCACCCTCG	180§	120
Vesicle trafficking protein (<i>unc-18</i>) F27D9.1a, intron 7	EE724186	unc18F unc18R	GCTAATGTAGCTATGGCTG GCATCCTCCATGATGTCC	429	250
Phosphoenolpyruvate carboxykinase (R11A5.4), intron 3	EE724187	pepckF pepckR	GGCTATGCGTCCATTTCATGGG GTCTTGC CGAACCAGTTGAC	246	128
Thin filament (F-actin) associated protein (<i>unc-87</i>) F08B6.4b, intron 2	EE724189	unc87F unc87R	GGAACTCCCAGGAACACCAGC CGTTCCTGACTGAAGCGGGAC	494	386
Calmodulin (<i>cmd-1</i>) T21H3.3, Intron 2	EE724190	cmd1F cmd1R	GGATACTGACAGTGAAGA CTCTCCAGATTCTGCAT	579	467
Casein kinase 1 (<i>kin-19</i>) C03C10.1, intron 1	EE724191	kin19F kin19R	CTTCAACTTTGCTCACGCAGG CCAATCATCTGATCTGCCAGC	320	255
Chitinase (T05H4.7), intron 2	EE724192	chitF chitR	CACATGCCAATCAAGAAACAGGGG CCAGTTGAAGAAATCCACCCTCG	257	150
Nucleolar protein (<i>lpd-7</i>) R13A5.12, intron 1	EE724193	lpd7F lpd7R	CGGTTAACTATATTTTCACGG GGATAGTACCACACAG	226	66
Glycosyl hydrolase (<i>cht-1</i> family 18) C04F6.3, intron 2	EE724194	cht1F cht1R	GGTTATGCTGAGATTATCGC GAATCGTCTTTCATCTTCCC	502	374
Serine/threonine protein phosphatase (<i>gsp-2</i>) F56C9.1, introns 1 and 2	EE724195	gsp2F gsp2R	CTGACAAGCTCAACTTGG GAAGGTTCATAGTATTGGCC	439†	124 108
Rab GDP dissociation inhibitor (<i>gdi-1</i>) Y57G11C.10, intron 1	EE724197	gdi1F gdi1R	CCGGGAACGAATGATGCG CTGTGCCAAGATCAGTCCGCTCG	471†	63 82
Hypothetical protein T19D7.5, introns 1 and 2	EE24198	hpF hpR	GATCACAGCCGAGGAGGCAGC CAGTGCAGGAGCGATGACCCG	381	245
Isocitrate lyase (<i>gei-7</i>) C05E4.9a, intron 4	EE724201	iclF iclR	CCGAAGGAAGTGAACAGATGCC GCATCATGTCTCAGCCACTGCC	342	169
Synaptosome-associated protein (<i>ric-4</i>) Y22F5A.3, introns 1, 2 and 3	EE724203	snap25F snap25R	GAGCTGAAGACTCTCAACTTGC GATTAATCGTGTCTAGAGCGCC	888‡	318 126 258
Regulator of G protein-signalling (<i>rgs-2</i>) F16H9.1a, intron 3	EE724204	rgs2F rgs2R	GCCTGGTCCCAATCCTTTGAC CCGCTTCAACTCTTCACAAGCC	199	70
Elongation factor-1 (<i>eft-3</i>) F31E3.5, intron 1	EE724216	eft3F eft3R	CATTGGACACGTCGATTCTGG GAACTCTCCTGTACCACATGC	405	71
G protein alpha subunit (<i>odr-3</i>) C34D1.3, intron 1	EU131131	odr3F odr3R	GGATCCAGCATCGTCAAGC GGGAATGTGTAGAATGTTGTCC	617§	470

The name of the homologous *Caenorhabditis elegans* sequence is given and, when available; the *C. elegans* gene name is also given in italics. †two introns present; ‡three introns present; §successful PCR but amplicons were not sequenced.

References

- Adams BJ, Fodor A, Koppenhöfer HS *et al.* (2006) Biodiversity and systematics of nematode-bacterium entomopathogens. *Biological Control*, **37**, 32–49.
- Akhurst RJ, Bedding RA (1986) Natural occurrence of insect pathogenic nematodes (Steinernematidae, Heterorhabditidae) in soil in Australia. *Journal of the Australian Entomology Society*, **25**, 241–244.
- Boemare NE, Akhurst RJ, Mourant RG (1993) DNA relatedness between *Xenorhabdus* spp. (Enterobacteriaceae), symbiotic bacteria of entomopathogenic nematodes, with a proposal to transfer *Xenorhabdus luminescens* to a new genus *Photorhabdus* gen. nov. *International Journal of Systematic Bacteriology*, **43**, 249–255.
- Dix I, Burnell AM, Griffin CT *et al.* (1992) The identification of biological species in the genus *Heterorhabditis* (Nematoda: Heterorhabditidae) by cross-breeding second generation amphimictic adults. *Parasitology*, **104**, 509–518.
- Graur D, Li WH (2000) *Fundamentals of Molecular Evolution*, 2nd edn. Sinauer Associates, Sunderland, Massachusetts.
- Palumbi SR, Baker CS (1994) Contrasting population structure from nuclear intron sequences and mtDNA of humpback whales. *Molecular Biology and Evolution*, **11**, 426–435.
- Thompson JD, Gibson TJ, Plewniak F *et al.* (1997) The Clustal_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research*, **25**, 4876–4882.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1 Cross-species comparison of size polymorphisms from 169 introns sequenced from eight species of the EPN genus *Heterorhabditis*. The size in base pairs for each intron is recorded (–, no amplification; **, intron absent) and the intron numbers refer to intron positions in *C. elegans*

Table S2 Gene identities and GenBank Accession numbers of the gDNA EPIC-PCR sequences obtained for *Heterorhabditis bacteriophora* and seven other *Heterorhabditis* species (–, EPIC-PCR product was not obtained)

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.